


Supplementary Data:**SOLUBILITY OF PHENYLBORONIC COMPOUNDS
IN WATER**

Paweł Leszczyński, Andrzej Sporzyński

Faculty of Chemistry, Warsaw University of Technology, Noakowskiego 3,
00-664 Warsaw, Poland*Mediterranean Journal of Chemistry* **2017**, 6(5), 200-207DOI: <http://dx.doi.org/10.13171/mjc65/01711030630-sporzynski>

Compound 1.

Method	$t/^{\circ}\text{C}$	x	Method	$t/^{\circ}\text{C}$	x	Method	$t/^{\circ}\text{C}$	x
A	31.7	0.00475	B	20.0	0.00370	C	25.0	0.00272
	35.5	0.00553		25.0	0.00421		30.0	0.00344
	38.7	0.00584		30.0	0.00465		35.0	0.00374
	40.5	0.00661		35.0	0.00521		40.0	0.00490
	45.0	0.00774		40.0	0.00614		45.0	0.00560
	48.4	0.00848		45.0	0.00665		50.0	0.00705
	53.6	0.00966		50.0	0.00848		60.0	0.00955

Compound 2.

Method	$t/^{\circ}\text{C}$	x	Method	$t/^{\circ}\text{C}$	x
A	34.6	0.00196	B	20.0	0.00141
	40.8	0.00226		25.0	0.00152
	42.5	0.00273		30.0	0.00168
	44.2	0.00278		35.0	0.00174
	50.2	0.00364		40.0	0.00189
	56.2	0.00398		45.0	0.00196
	65.9	0.00458		50.0	0.00207

Compound 3.

Method	$t/^{\circ}\text{C}$	x	Method	$t/^{\circ}\text{C}$	x
B	30.0	0.00008	C	20.0	0.00004
	35.0	0.00009		25.0	0.00007
	40.0	0.00010		30.0	0.00007
	45.0	0.00012		35.0	0.00005
	50.0	0.00011		40.0	0.00013
	60.0	0.00023		45.0	0.00006
	65.0	0.00029		50.0	0.00009
				55.0	0.00010
				60.0	0.00014

Compound 4.

Method	$t/^{\circ}\text{C}$	x
C	20.0	0.00014
	25.0	0.00028
	30.0	0.00018
	35.0	0.00035
	40.0	0.00023
	45.0	0.00053
	50.0	0.00049
	55.0	0.00059
	60.0	0.00084
	65.0	0.00096
	70.0	0.00148

Compound 5.

Method	$t/^{\circ}\text{C}$	x
C	30.0	0.00151
	40.0	0.00175
	50.0	0.00223
	55.0	0.00360
	60.0	0.00437

Compound 6.

Method	$t/^{\circ}\text{C}$	x
C	30.0	0.00017
	40.0	0.00023
	50.0	0.00024
	55.0	0.00034
	60.0	0.00033